

## Ramapo College Immunization Requirements

#### **DUE DATE**

Fall Semester Start: July 31st

Spring Semester Start: January 2<sup>nd</sup>

### **IMMUNIZATION POLICY**

Failure to complete health requirements will result in a registration hold.

All registered students are required to submit this form if you are taking classes (in person, online, or virtual).

### REQUIRED IMMUNIZATION FORMS

1. Take this packet to your health ca

# **IMMUNIZATION RECORD**

PART 1: COMPLETED BY THE STUDENT. All information must be printed legibly or form cannot be processed.				
Last Name:	First Name:	Middle Initial:		
DOB:	Start Date:Fall Spring Year:	Cell Number:		
Full	•	•		

Full-



### **TOOL FOR INSTITUTIONAL USE-ATTACHMENT 1**

### Part I: Tuberculosis (TB) Screening Questionnaire (to be completed by incoming students)

Please answer the following questions (1 through 6):

1. Have you ever had close contact with persons known or suspected to have active TB disease? Yes No

2. Were you born in one of the countries or territories listed below that have a high incidence of active TB disease? (If yes, please CIRCLE the country below)

Yes

No

Afghanistan

Algeria

Angola

Anguilla

Argentina

Armenia

Azerbaijan

Bangladesh

Belarus

Belize

Benin Bhutan

Bolivia (Plurinational State

of

Bosnia and Herzegovina

Botswana

Brazil

Brunei Darussalam



3.	INTERFERON GAMMA RELEASE ASSAY (IGRA): MAY BE COMPLETED OUTSIDE OF THE UNITED STATES BUT LAB REPORT IS REQUIRED IN ENGLISH.		
	Date Obtained:/ (circle method) QFT-GIT T-Spot other		
	Result: negative positive indeterminate borderline (T-Spot only)		
4.	CHEST X-RAY: REQUIRED IF TST OR IGRA IS POSITIVE AND MUST BE PERFORMED IN THE UNITED STATES. LAB REPORT IS REQUIRED IN ENGLISH.  Date of chest X-ray:/ Result: normal abnormal		

### Part III: Management of Positive TST or IGRA

All students with a positive TST or IGRA with no signs of active disease on chest X-ray should receive a recommendation to be treated for latent TB with appropriate medication. However, students in the following groups are at increased risk of progressions from LTBI to TB disease and should be prioritized to begin treatment as soon as possible.

Infected with HIV