# IMMUNIZATION RECORD

| **PART 1: COMPLETED BY THE STUDENT. All information must be printed legibly or form cannot be processed.** | | |
|---|---|---|
| Last Name: | First Name: | Middle Initial: |
| DOB: | Start Date: ___Fall ____ Spring Year: | Cell Number: |

Full-

**Part II: Clinical Assessment to be completed by Health Care Provider**

Clinicians should review and verify the information in Part I. Persons who answered NO to all questions in Part I do not need further testing. Persons who answered YES to any of the questions in Part I are candidates for either the Mantoux tuberculin skin test (TST) or the Interferon Gamma Release Assay (IGRA), unless a previous positive test has been documented.

History of a positive TB skin test or IGRA blood test? (If yes, document below.) ☐ Yes ☐ No

History of BCG vaccination? (If yes, consider IGRA if possible.) ☐ Yes ☐ No

**1. TB SYMPTOM CHECK**

Does the student have signs or symptoms of active pulmonary tuberculosis disease? ☐ Yes ☐ No

If No, proceed to 2 or 3.

**If Yes, check below:**

- o Cough (especially if lasting for 3 weeks or longer) with or without sputum production
- o Coughing up blood (hemoptysis)
- o Chest pain
- o Loss of appetite
- o Unexplained weight loss
- o Night sweats
- o Fever

Proceed with additional evaluation to exclude active tuberculosis disease, including tuberculin skin testing, chest X-ray, and sputum evaluation as indicated.

**2. TUBERCULIN SKIN TEST (TST): MUST BE PERFORMED IN THE UNITED STATES (IF CURRENTLY LIVING OUTSIDE OF THE UNITED STATES, GO TO #3).**

(TST result should be recorded as actual millimeters [mm] of induration, transverse diameter; if no induration, write ñ0.òThe TST interpretation should be based on mm of induration as well as risk factors.)**

Date Given:____/_____/____ Date Read: ____/_____/____

Result: _____mm of induration **Interpretation: positive____ negative____

****INTERPRETATION GUIDELINES**

**>5 mm is positive:**

Recent close contacts of an individual with infectious TB

Persons with fibrotic changes on a prior chest X-ray, consistent with past TB disease

Organ transplant recipients and other immunosuppressed persons (including receiving equivalent of >15mg/d of prednisone for >1 month)

HIV-infected persons

**> 10 mm is positive:**

Recent arrivals to the U.S. (<5 years) from high-prevalence areas or who resided in one for a significant amount of time (The significance of the travel exposure should be discussed with a health care provider and evaluated.)

Injection drug users

Mycobacteriology laboratory personnel

Residents, employees, or volunteers in high-risk congregate settings

Persons with medical conditions that increase the risk of progression to TB disease, including silicosis, diabetes mellitus, chronic renal failure, certain types of cancer (leukemias and lymphomas, cancers of the head, neck, or lung), gastrectomy or jejunoileal bypass and weight loss of at least 10% below ideal body weight.

**>15 mm is positive:**

Persons with no known risk factors for TB who, except for certain testing programs required by law or regulation, would otherwise not be tested.